

Section 5.2: The Characteristic Equation

Recall

- ▶ For $n \times n$ matrix A , a nonzero vector \mathbf{x} and scalar λ are an **eigenvector–eigenvalue** pair provided $A\mathbf{x} = \lambda\mathbf{x}$.
- ▶ For eigenvalue λ , the subspace $\{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\} = \text{Nul}(A - \lambda I)$ is called the **eigenspace** of A corresponding to λ .
- ▶ The polynomial equation $\det(A - \lambda I) = 0$ is called the **characteristic equation**, and λ is an eigenvalue if and only if it's a solution to this equation.

Recall

- ▶ The eigenvalues of a triangular matrix are the diagonal entries.
- ▶ A matrix A is singular if and only if $\lambda = 0$ is an eigenvalue (i.e., it's invertible if and only if $\lambda = 0$ is NOT an eigenvalue).
- ▶ The **algebraic multiplicity** of an eigenvalue is its multiplicity as a root of the characteristic equation.
- ▶ The **geometric multiplicity** of an eigenvalue is the dimension of its corresponding eigenspace.

Similarity

Definition:

Two $n \times n$ matrices A and B are said to be **similar** if there exists an invertible matrix P such that

$$B = P^{-1}AP.$$

$$A = PBP^{-1}$$

The mapping $A \mapsto P^{-1}AP$ is called a **similarity transformation**^a.

^a**Note:** similarity is NOT related to row equivalence.

Theorem:

If A and B are similar matrices, then they have the same characteristic equation, and hence the same eigenvalues.

Example

Show that $A = \begin{bmatrix} -18 & 42 \\ -7 & 17 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & 0 \\ 0 & -4 \end{bmatrix}$ are similar with the matrix P for the similarity transformation given by $P = \begin{bmatrix} 2 & 3 \\ 1 & 1 \end{bmatrix}$.

We'll show that $B = P^{-1}AP$.

We need P^{-1} . $\det(P) = 2(1) - 3(1) = -1$

$$P^{-1} = \frac{1}{-1} \begin{bmatrix} 1 & -3 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} -1 & 3 \\ 1 & -2 \end{bmatrix}$$

$$P^{-1}AP = \begin{bmatrix} -1 & 3 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} -18 & 42 \\ -7 & 17 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 3 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} 6 & -12 \\ 3 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & 0 \\ 0 & -4 \end{bmatrix}$$

Example Continued...

Show that the columns of P are eigenvectors of A where

$$A = \begin{bmatrix} -18 & 42 \\ -7 & 17 \end{bmatrix} \quad \text{and} \quad P = \begin{bmatrix} 2 & 3 \\ 1 & 1 \end{bmatrix}.$$

$$= [\vec{p}_1 \quad \vec{p}_2]$$

We need to show that $A\vec{p}_1 = \lambda_1\vec{p}_1$ and

$A\vec{p}_2 = \lambda_2\vec{p}_2$ for some λ_1, λ_2 .

$$A\vec{p}_1 = \begin{bmatrix} -18 & 42 \\ -7 & 17 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A\vec{p}_2 = \begin{bmatrix} -18 & 42 \\ -7 & 17 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} -12 \\ -4 \end{bmatrix} = -4 \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

Eigenvalues of a real matrix need not be real

Find the eigenvalues of the matrix $A = \begin{bmatrix} 4 & 3 \\ -5 & 2 \end{bmatrix}$.

Solve $\det(A - \lambda I) = 0$

$$\det(A - \lambda I) = \det \begin{bmatrix} 4 - \lambda & 3 \\ -5 & 2 - \lambda \end{bmatrix} = (4 - \lambda)(2 - \lambda) + 15$$

$$= \lambda^2 - 6\lambda + 8 + 15$$

$$= \lambda^2 - 6\lambda + 23$$

Characteristic eqn is

$$\lambda^2 - 6\lambda + 23 = 0$$

Using the quadratic formula

$$\lambda = \frac{6 \pm \sqrt{(-6)^2 - 4(1)(23)}}{2(1)}$$

$$= \frac{6 \pm \sqrt{36 - 92}}{2} = \frac{6 \pm \sqrt{-56}}{2}$$

$$= \frac{6 \pm 2\sqrt{14}i}{2} = 3 \pm \sqrt{14}i$$

A has no eigenvectors in \mathbb{R}^2 .

Section 5.3: Diagonalization

Motivational Example:

Determine the eigenvalues of the matrix D^6 (that's D raised to the sixth power), where $D = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix}$.

Let's find D^6 .

$$D^2 = D D = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 2^2 & 0 \\ 0 & (-1)^2 \end{bmatrix}$$

$$D^3 = D^2 D = \begin{bmatrix} 2^2 & 0 \\ 0 & (-1)^2 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 2^3 & 0 \\ 0 & (-1)^3 \end{bmatrix}$$

$$D^6 = \begin{bmatrix} 2^6 & 0 \\ 0 & (-1)^6 \end{bmatrix}$$

The eigenvalues of D^6 are

$$\lambda_1 = 2^6 \quad \text{and} \quad \lambda_2 = (-1)^6$$

Recall that a matrix D is diagonal if it is both upper and lower triangular (its only nonzero entries are on the diagonal).

Theorem

If D is a diagonal matrix with diagonal entries d_{ij} , then D^k is diagonal with diagonal entries d_{ij}^k for positive integer k . Moreover, the eigenvalues of D are the diagonal entries.

$$D = \begin{bmatrix} d_{11} & 0 & \cdots & 0 \\ 0 & d_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn} \end{bmatrix} \implies D^k = \begin{bmatrix} d_{11}^k & 0 & \cdots & 0 \\ 0 & d_{22}^k & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn}^k \end{bmatrix}$$

Powers and Similarity

Suppose A and B are similar matrices with similarity transform matrix P —i.e., $B = P^{-1}AP$. Show that

- A^2 and B^2 are similar with the same P ,
- A^3 and B^3 are similar with the same P .

$$\text{Given } B = P^{-1}AP$$

$$\begin{aligned} B^2 &= B B \\ &= (P^{-1}AP)(P^{-1}AP) \\ &= P^{-1}A \underbrace{(PP^{-1})}_I A P \\ &= P^{-1}A I A P \end{aligned}$$

$$= P^{-1} A^2 P$$

$$B^3 = B^2 B$$

$$= (P^{-1} A^2 P)(P^{-1} A P)$$

$$= P^{-1} A^2 (P P^{-1}) A P$$

$$= P^{-1} A^2 A P$$

$$= P^{-1} A^3 P$$

$$B^n = (P^{-1} A P)(P^{-1} A P)(P^{-1} A P) \dots (P^{-1} A P)$$

$$= P^{-1} A^n P$$

Diagonalizability

Defintion:

An $n \times n$ matrix A is called **diagonalizable** if it is similar to a diagonal matrix D . That is, provided there exists a nonsingular matrix P such that $D = P^{-1}AP$ —i.e. $A = PDP^{-1}$.

Theorem:

The $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors. In this case, the matrix P is the matrix whose columns are the n linearly independent eigenvectors of A .

Example

Diagonalize the matrix A if possible. $A = \begin{bmatrix} 1 & 3 & 3 \\ -3 & -5 & -3 \\ 3 & 3 & 1 \end{bmatrix}$

- Find eigenvalues
- Find eigenvectors
- If we get n lin. independent eigenvectors, form P and find $D = P^{-1}AP$.

Find the eigenvalues.

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 & 3 \\ -3 & -5 - \lambda & -3 \\ 3 & 3 & 1 - \lambda \end{bmatrix}$$

$$= (1-\lambda) \begin{vmatrix} -s-\lambda & -3 \\ 3 & 1-\lambda \end{vmatrix} - 3 \begin{vmatrix} -3 & -3 \\ 3 & 1-\lambda \end{vmatrix} + 3 \begin{vmatrix} -3 & -s-\lambda \\ 3 & 3 \end{vmatrix}$$

$$= (1-\lambda) \left(-(s+\lambda)(1-\lambda) + 9 \right) - 3 \left(-3(1-\lambda) + 9 \right) + 3 \left(-9 + 3(s+\lambda) \right)$$

$$= (1-\lambda) \left(-(-\lambda^2 - 4\lambda + 5) + 9 \right) - 3(3\lambda + 6) + 3(3\lambda + 6)$$

$$= (1-\lambda)(\lambda^2 + 4\lambda + 4)$$

$$= (1-\lambda)(\lambda+2)^2$$

The characteristic eqn is

$$(1-\lambda)(\lambda+2)^2 = 0$$

The eigenvalues are $\lambda_1 = 1$ and $\lambda_2 = \lambda_3 = -2$

Find e.vectors. For $\lambda_1 = 1$

$$A - 1I = \begin{bmatrix} 0 & 3 & 3 \\ -3 & -6 & -3 \\ 3 & 3 & 0 \end{bmatrix} \quad \text{we need to solve } (A - 1E)\vec{x} = \vec{0}$$

$$\begin{array}{l} \text{rref} \\ \rightarrow \end{array} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{array}{l} x_1 = x_3 \\ x_2 = -x_3 \\ \text{nd} \\ x_3 \text{ is free} \end{array}$$

An eigenvector is $\vec{x} = x_3 \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$

Let $\vec{v}_1 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$.

$$\text{For } \lambda_{2,3} = -2, \quad A - (-2)I = \begin{bmatrix} 3 & 3 & 3 \\ -3 & -3 & -3 \\ 3 & 3 & 3 \end{bmatrix}$$

$$(A + 2I)\vec{x} = \vec{0}$$

$$A + 2I \xrightarrow{\text{rref}} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$x_1 = -x_2 - x_3$$

x_2, x_3 free

$$\vec{x} = x_2 \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

$$\text{Let } \vec{v}_2 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \quad \vec{v}_3 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

$$\text{Let } P = [\vec{v}_1 \quad \vec{v}_2 \quad \vec{v}_3] = \begin{bmatrix} 1 & -1 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$$P^{-1} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ -1 & -1 & 0 \end{bmatrix}$$

$$D = P^{-1}AP = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -2 \end{bmatrix}$$